

MIT Open Access Articles

p-adic dimensions in symmetric
tensor categories in characteristic *p*

The MIT Faculty has made this article openly available. **Please share** how this access benefits you. Your story matters.

Citation: Etingof, Pavel, Nate Harman, and Victor Ostrik, "p-adic dimensions in symmetric tensor categories in characteristic p." Quantum topology 9, 1 (Feb. 2018): p. 119-40 doi:: 10.4171/QT/104 ©2018 Author(s)

As Published: 10.4171/QT/104

Publisher: European Mathematical Publishing House

Persistent URL: <https://hdl.handle.net/1721.1/124324>

Version: Author's final manuscript: final author's manuscript post peer review, without publisher's formatting or copy editing

Terms of use: Creative Commons Attribution-Noncommercial-Share Alike



**p -ADIC DIMENSIONS IN SYMMETRIC TENSOR
CATEGORIES IN CHARACTERISTIC p**

PAVEL ETINGOF, NATE HARMAN, VICTOR OSTRIK

ABSTRACT. To every object X of a symmetric tensor category over a field of characteristic $p > 0$ we attach p -adic integers $\mathrm{Dim}_+(X)$ and $\mathrm{Dim}_-(X)$ whose reduction modulo p is the categorical dimension $\dim(X)$ of X , coinciding with the usual dimension when X is a vector space. We study properties of $\mathrm{Dim}_\pm(X)$, and in particular show that they don't always coincide with each other, and can take any value in \mathbb{Z}_p . We also discuss the connection of p -adic dimensions with the theory of λ -rings and Brauer characters.

1. INTRODUCTION

Let \mathbf{k} be an algebraically closed field of characteristic $p > 0$, and let \mathcal{C} be a symmetric tensor (in particular, abelian) category over \mathbf{k} . Let $X \in \mathcal{C}$. Then to X one can attach its categorical dimension $\dim(X) \in \mathbf{k}$, and it is shown in [EGNO], Exercise 9.9.9(ii) that in fact $\dim(X) \in \mathbb{F}_p \subset \mathbf{k}$. The main result of the paper is that $\dim(X)$ is the reduction modulo p of a richer invariant of X , which takes values in \mathbb{Z}_p — the p -adic dimension of X . In fact, we define two kinds of p -adic dimensions — the symmetric p -adic dimension $\mathrm{Dim}_+(X)$ and the exterior p -adic dimension $\mathrm{Dim}_-(X)$, which for vector spaces coincide with the usual dimension of X .

Specifically, recall that if $t = \dots t_2 t_1 t_0 \in \mathbb{Z}_p$ is a p -adic integer with digits t_0, t_1, t_2, \dots (i.e., $t = \sum_{j \geq 0} t_j p^j$), then we can define the power series

$$(1+z)^t := \prod_{j=0}^{\infty} (1+z^{p^j})^{t_j} \in \mathbb{F}_p[[z]],$$

and t is uniquely determined by the series $(1+z)^t$. This series has all the properties of the exponential function, i.e., $(1+z)^{t+s} = (1+z)^t(1+z)^s$, $((1+z)^t)^s = (1+z)^{ts}$, etc.

Now, the p -adic dimensions of X are defined by the formulas

$$(1) \quad \sum_{j=0}^{\infty} \dim(S^j X) z^j = (1-z)^{-\mathrm{Dim}_+(X)},$$

and

$$(2) \quad \sum_{j=0}^{\infty} \dim(\wedge^j X) z^j = (1+z)^{\text{Dim}_-(X)},$$

where $S^i X$ and $\wedge^i X$ are the symmetric and exterior powers of X . The main challenge is to show that $\text{Dim}_{\pm}(X)$ exist. We do it in Section 2.

It follows immediately from the definition that

$$\text{Dim}_{\pm}(X \oplus Y) = \text{Dim}_{\pm}(X) + \text{Dim}_{\pm}(Y)$$

and that p -adic dimension is functorial, i.e. for any (exact) symmetric tensor functor F we have $\text{Dim}_{\pm}(F(X)) = \text{Dim}_{\pm}(X)$. Also, using Deligne’s categories in characteristic p , we show that $\text{Dim}_+(X)$ and $\text{Dim}_-(X)$ can take any value in \mathbb{Z}_p . On the other hand, we show that in general

$$\text{Dim}_+(X) \neq \text{Dim}_-(X), \text{Dim}_{\pm}(X \otimes Y) \neq \text{Dim}_{\pm}(X)\text{Dim}_{\pm}(Y),$$

and at least in characteristic 2 the dimensions Dim_{\pm} are not always additive on exact sequences, although these properties do hold for “good” objects.

We will discuss applications of p -adic dimensions in subsequent papers. One of the motivations is to formulate the universal property of Deligne’s categories $\text{Rep}(S_t)$ and $\text{Rep}(GL_t)$ in positive characteristic, defined by P. Deligne in his letter to the third author, [De3] (see [Ha], Section 3.3, for a summary). Namely, for any nontrivial ultrafilter \mathcal{U} on the set of natural numbers Deligne constructed the tensor categories of superexponential growth $\text{Rep}(S_{\mathcal{U}})$, $\text{Rep}(GL_{\mathcal{U}})$ over \mathbf{k} , analogous to Deligne’s categories $\text{Rep}(S_t)$, $\text{Rep}(GL_t)$ in characteristic zero (see [DM, De1, De2] and [EGNO], Section 9.12). These categories are obtained by “ p -adic interpolation” of the ordinary representation categories of S_n and GL_n along \mathcal{U} . Deligne conjectured that they depend only on the p -adic integer $t := \lim_{\mathcal{U}} n$ (so that one may denote them by $\text{Rep}(S_t)$, $\text{Rep}(GL_t)$), and for $\text{Rep}(S_{\mathcal{U}})$ this has now been proved in [Ha] for $p \geq 5$. In characteristic zero, the categories $\text{Rep}(S_t)$ and $\text{Rep}(GL_t)$ have nice universal properties; for instance, symmetric tensor functors from $\text{Rep}(GL_t)$, $\text{Rep}(S_t)$ to a symmetric tensor category \mathcal{C} correspond to objects (respectively, commutative Frobenius algebras) X in \mathcal{C} of dimension t such that for any partition λ , the Schur functor $\mathcal{S}^{\lambda} X$ is nonzero. The notion of p -adic dimension should allow one to generalize these properties to positive characteristic; namely, one should expect that symmetric tensor functors from $\text{Rep}(GL_t)$, $\text{Rep}(S_t)$

to \mathcal{C} correspond to “good” objects (respectively, commutative Frobenius algebras) X in \mathcal{C} of p -adic dimension t . We plan to discuss these universal properties more precisely elsewhere.

The organization of the paper is as follows. In Section 2 we define the p -adic dimensions and prove their existence. In Section 3 we study the properties of p -adic dimensions, and discuss their connection with λ -rings and Brauer characters.

Acknowledgements. We are very grateful to Pierre Deligne for his letter to V.O., which led to this work. We thank Haynes Miller for Remark 3.4 and Siddharth Venkatesh for useful discussions. The work of P.E. and N.H. was partially supported by the NSF grant DMS-1502244. The work of N. H. was also partially supported by the National Science Foundation Graduate Research Fellowship under Grant No. 1122374.

2. p -ADIC DIMENSIONS

2.1. Preliminaries on tensor categories. Let \mathbf{k} be an algebraically closed field of characteristic $p > 0$, and \mathcal{C} a symmetric tensor category over \mathbf{k} (see [EGNO], Definitions 4.1.1, 8.1.2). For $X, Y \in \mathcal{C}$, we have the commutativity constraint $c_{X,Y} : X \otimes Y \rightarrow Y \otimes X$. Since $c_{Y,X} \circ c_{X,Y} = 1_{X \otimes Y}$, we have a natural action of the symmetric group S_n on $X^{\otimes n}$ defined by $s_{i,i+1} \mapsto c_i := 1^{\otimes i-1} \otimes c_{X,X} \otimes 1^{n-i-1}$, where $s_{i,j} \in S_n$ is the transposition of i and j .

Recall from [EGNO] that to every endomorphism $A : X \rightarrow X$ of an object $X \in \mathcal{C}$ we can attach its trace

$$\mathrm{Tr}_X(A) := \mathrm{ev}_X \circ c_{X,X^*} \circ (A \otimes 1) \circ \mathrm{coev}_X \in \mathrm{End}(\mathbf{1}) = \mathbf{k},$$

and in particular we can define the dimension of X by $\dim(X) = \mathrm{Tr}_X(1) \in \mathbf{k}$. We have $\mathrm{Tr}_{X^*}(A^*) = \mathrm{Tr}_X(A)$, hence $\dim(X^*) = \dim(X)$.

Lemma 2.1. *Let $\sigma \in S_n$ be a permutation. Then $\mathrm{Tr}_{X^{\otimes n}}(\sigma) = \dim(X)^{c(\sigma)}$, where $c(\sigma)$ is the number of cycles of σ .*

Proof. This is easy to prove diagrammatically. Consider the flat braid b_σ corresponding to σ . Then the trace of σ is depicted by the flat link obtained by gluing the i -th input of b_σ to its i -th output for all $i \in [1, n]$. It is easy to see that this flat link has $c(\sigma)$ components, and each component contributes a factor of $\dim(X)$, which implies the statement. \square

Lemma 2.2. ([EGNO], Exercise 9.9.9(ii)) *One has $\dim(X) \in \mathbb{F}_p$.*

Proof. Let s be the cyclic permutation acting on $X^{\otimes p}$. Then $s^p = 1$, so $(1 - s)^p = 0$ (as we are in characteristic p), and hence $1 - s$ is

nilpotent. Thus $\text{Tr}_{X^{\otimes p}}(1-s) = 0$. But by Lemma 2.1, this trace equals $\dim(X)^p - \dim(X)$. Thus $\dim(X)^p = \dim(X)$, so $\dim(X) \in \mathbb{F}_p$. \square

Now recall the definition of symmetric and exterior powers of X . This definition is given in [EGNO], Section 9.9, under the assumption of characteristic zero; the discussion also applies to characteristic $p > 2$, but some changes are needed for $p = 2$.

Let $\Lambda^2 X$ be the image of $c_{X,X} - 1$ in $X \otimes X$. We define the symmetric algebra SX to be the quotient of the tensor algebra TX by the two-sided ideal generated by $\Lambda^2 X$. Then SX is a \mathbb{Z}_+ -graded algebra: $SX = \bigoplus_{n \geq 0} S^n X$. The object $S^n X$ is the object of coinvariants of S_n in $X^{\otimes n}$, and is called *the n -th symmetric power of X* . We also define the *dual n -th symmetric power* $\mathbb{S}^n X := (S^n X^*)^*$. We have a natural inclusion $\mathbb{S}^n X \subset X^{\otimes n}$ as the subobject of S_n -invariants; i.e., $\mathbb{S}^n X$ is the intersection of the kernels of $c_i - 1$ for $i = 1, \dots, n-1$. We also have a natural projection $X^{\otimes n} \rightarrow S^n X$. The composition of these two maps is a morphism $\phi_n : \mathbb{S}^n X \rightarrow S^n X$. This morphism is an isomorphism for $n < p$ but not for $n \geq p$.

Similarly, consider $\mathbb{S}^2 X \subset X \otimes X$, and define the exterior algebra $\wedge X$ as the quotient of TX by the two-sided ideal generated by $\mathbb{S}^2 X$. Then $\wedge X = \bigoplus_{n \geq 0} \wedge^n X$, and $\wedge^n X$ is a quotient of $X^{\otimes n}$ called the *n -th exterior power of X* . We also define the *dual n -th exterior power* $\Lambda^n X := (\wedge^n X^*)^* \subset X^{\otimes n}$, which is the intersection of the images of $c_i - 1$ for $i = 1, \dots, n-1$ (so for $n = 2$ we recover $\Lambda^2 X$ defined above). We have a morphism $\psi_n : \Lambda^n X \rightarrow \wedge^n X$ which is an isomorphism for $n < p$ but not for $n \geq p$.

If $p > 2$ then $S^2 X \cong \mathbb{S}^2 X$, so the exterior power $\wedge^n X$ can be defined as the object of anticoinvariants of S_n in $X^{\otimes n}$, while the dual exterior power $\Lambda^n X \subset X^{\otimes n}$ is the subobject of antiinvariants (the intersection of the kernels of $c_i + 1$ or, equivalently, images of $c_i - 1$ over all i).

Also, in any characteristic these definitions of symmetric and exterior algebras and powers coincide with the usual ones if X is a vector space over \mathbf{k} .

Finally, note that $\dim(S^i X) = \dim(\mathbb{S}^i X^*)$ and $\dim(\wedge^i X) = \dim(\Lambda^i X^*)$.

2.2. The p -adic dimensions. Let $X \in \mathcal{C}$. The following theorem is our main result.

Theorem 2.3. (i) *There exists a unique $d_+ = d_+(X) \in \mathbb{Z}_p$ such that*

$$\sum_{j \geq 0} \dim(\mathbb{S}^j X) z^j = (1-z)^{-d_+}.$$

(ii) There exists a unique $d_- = d_-(X) \in \mathbb{Z}_p$ such that

$$\sum_{j \geq 0} \dim(\Lambda^j X) z^j = (1 + z)^{d_-}.$$

Proof. (i) Let $d_r := \dim(\mathbb{S}^r X)$. Let n be a positive integer, and $\lambda = (\lambda_1, \lambda_2, \dots, \lambda_m)$ be a partition of n of length m . Let

$$S_\lambda := S_{\lambda_1} \times \dots \times S_{\lambda_m} \subset S_n$$

be the Young subgroup of λ . Let

$$\mathbb{S}^\lambda X := \mathbb{S}^{\lambda_1} X \otimes \dots \otimes \mathbb{S}^{\lambda_m} X = (X^{\otimes n})^{S_\lambda}.$$

Given a double coset $B \in S_\lambda \backslash S_n / S_\lambda$, we have a canonical endomorphism $A_B : \mathbb{S}^\lambda X \rightarrow \mathbb{S}^\lambda X$. Namely, write B as a disjoint union of left cosets B_i of S_λ in S_n , and pick elements $b_i \in B_i$. Then define the endomorphism A' of $X^{\otimes n}$ by $A' := \sum_i b_i$.

Claim 1. A' restricts to an endomorphism A_B of $\mathbb{S}^\lambda X$, and the resulting endomorphism is independent on the choice of the b_i .

Indeed, if $b'_i \in B_i$ is another choice, then $b'_i = b_i g_i$, where $g_i \in S_\lambda$, so $b'_i = b_i$ on $\mathbb{S}^\lambda X$. Also, for any $g \in S_\lambda$ we have $g \sum_i b_i = \sum_i g b_i$, and $g b_i$ is a representative of the coset $g B_i$. But the union of $g B_i$ over all i is disjoint and equals B , so $\sum_i g b_i = \sum_i b_i$ on $\mathbb{S}^\lambda X$. This proves the claim.

Lemma 2.4. *There exists a universal polynomial $P_B(x_1, x_2, \dots, x_{\lambda_1})$ with integer coefficients such that for a tensor category \mathcal{C} over a field of any characteristic and any $X \in \mathcal{C}$ we have $\text{Tr}_{\mathbb{S}^\lambda X}(A_B) = P_B(d_1, d_2, \dots, d_{\lambda_1})$.*

Proof. The proof is by reverse induction in the length m of λ , starting with length n . If $\text{length}(\lambda) = n$ then $\mathbb{S}^\lambda X = X^{\otimes n}$, $S_\lambda = 1$, $B \in S_n$, $A_B = B$, and $\text{Tr}_{\mathbb{S}^\lambda X}(A_B) = d_1^{c(B)}$ by Lemma 2.1, which gives the base of induction. So assume that $\text{length}(\lambda) = m < n$ and the statement is known for all larger lengths. If B is the double coset of 1 then $A_B = 1$, and $\text{Tr}_{\mathbb{S}^\lambda X}(A_B) = \prod_{i=1}^m d_{\lambda_i}$, as required. So we may assume without loss of generality that B is the double coset of $g \in S_n$, where $g \notin S_\lambda$. Let $H \subset S_\lambda$ be the intersection of S_λ with $g S_\lambda g^{-1}$. Recall that double cosets of S_λ in S_n are labeled by m by m matrices $\mathbf{a} = (a_{ij})$ of nonnegative integers such that $\sum_i a_{ij} = \lambda_i$ and $\sum_j a_{ij} = \lambda_j$ (see [St], Exercise 7.77). By choosing g to be of minimal length, we may assume that

$$g^{-1} H g = S_{\mathbf{a}} := \prod_{i=1}^m \prod_{j=1}^m S_{a_{ij}} \subset S_\lambda,$$

where (a_{ij}) is a partition of λ_i (nontrivial for at least one i); see [S], Lemma 2. Let c_s be representatives of elements s of S_λ/H in S_λ . Then we can write A_B as

$$A_B = \sum_{s \in S_\lambda/H} c_s g$$

(note that this is independent on the choice of c_s since $g^{-1}Hg \subset S_\lambda$).

Let $\mathbb{S}^{\mathbf{a}}X := \otimes_i \otimes_j \mathbb{S}^{a_{ij}}X$, and let $A' : \mathbb{S}^{\mathbf{a}}X \rightarrow X^{\otimes n}$ be defined by the same formula as A_B , i.e.,

$$A' = \sum_{s \in S_\lambda/H} c_s g.$$

Claim 2. A' lands in $\mathbb{S}^\lambda X \subset X^{\otimes n}$.

Indeed, let $y \in S_\lambda$. Then

$$yA' = \sum_s y c_s g = \sum_s c_s b_s g = \sum_s c_s g b'_s,$$

where $b_s \in H$ and $b'_s = g^{-1}b_s g \in g^{-1}Hg = S_{\mathbf{a}}$. So b'_s acts trivially on $\mathbb{S}^{\mathbf{a}}X$, hence $yA' = A'$, and thus A' lands in $\mathbb{S}^\lambda X$, as claimed.

It is clear that A' restricts on $\mathbb{S}^\lambda X$ to A_B . Thus, $\mathrm{Tr}_{\mathbb{S}^\lambda X}(A_B) = \mathrm{Tr}_{\mathbb{S}^{\mathbf{a}}X}(A')$. But $\mathrm{Tr}_{\mathbb{S}^{\mathbf{a}}X}(A')$ is given by some universal polynomial in $d_1, \dots, d_{\lambda_1}$ with integer coefficients by the induction assumption, as A' is a sum of endomorphisms of the form $A_{B'}$, where B' is a double coset of $S_{\mathbf{a}}$ in S_n , and $S_{\mathbf{a}}$ is a parabolic subgroup corresponding to the partition \mathbf{a} with $\mathrm{length}(\mathbf{a}) > m$. This proves the lemma. \square

Lemma 2.5. *Suppose that the ground field has characteristic $p > 0$. Then for each n there exists a universal polynomial $Q_n(z_0, z_1, \dots)$ over \mathbb{F}_p such that $d_n = Q_n(d_1, d_p, d_{p^2}, \dots)$. Moreover, Q_n is unique if we require that it is of degree $< p$ in every variable. This unique polynomial is given by the formula*

$$(3) \quad Q_n(z_0, z_1, z_2, \dots) = (-1)^n \prod_{i=0}^k \binom{(-1)^{p^i} z_i}{n_i},$$

where n_i are the digits of the base p expansion of n .

Proof. To prove existence, use induction in n . The base $n = 1$ is trivial. We may assume that n is not a power of p (otherwise, if $n = p^i$, then $Q_n(z) = z_i$). Suppose the statement is known below n . Let p^i be the largest power of p dividing n ($i \geq 0$). Then it is easy to see that the binomial coefficient $\binom{n}{p^i}$ is not zero in \mathbb{F}_p . Therefore, $\mathbb{S}^n X$ is the image

of the following projector P acting on $\mathbb{S}^{n-p^i} X \otimes \mathbb{S}^{p^i} X$:

$$P = \binom{n}{p^i}^{-1} \sum_s c_s,$$

where c_s are representatives of the elements $s \in S_n/S_{n-p^i} \times S_{p^i}$ in S_n . Thus, $d_n = \text{Tr}(P)$. But by Lemma 2.4, $\text{Tr}(P)$ is given by a universal polynomial over \mathbb{F}_p in d_1, \dots, d_r , where

$$r = \max(n - p^i, p^i).$$

Since $r < n$, by the induction assumption, d_i , $i \leq r$ are universal polynomials over \mathbb{F}_p of d_1, d_p, d_{p^2}, \dots . Hence so is d_n . This shows that the polynomial Q_n exists.

Moreover, since by Lemma 2.2 we have $d_i \in \mathbb{F}_p$, in Q_n we can replace z_i^p with z_i , and thus can choose Q_n in such a way that it has degree $< p$ with respect to each variable.

Now we want to show that Q_n with this degree requirement is unique and is given by formula (3). To do so, pick some Q_n satisfying the degree requirement, and let

$$f(z_0, \dots, z_k) = Q_n(z_0, \dots, z_k) - (-1)^n \prod_{i=0}^k \binom{(-1)^{p^i} z_i}{n_i},$$

for sufficiently large k . Let $N_0, N_1, \dots, N_k \in [0, \dots, p-1]$, and let

$$N = p^{k+1} - \sum_{i=0}^k N_i p^i.$$

Let \mathcal{C} be the category of vector spaces, and $X = \mathbf{k}^N$. Then $d_r = \binom{N+r-1}{r} = (-1)^r \binom{-N}{r}$. By Lucas' theorem, for $r < p^{k+1}$ we then get

$$d_r = (-1)^r \prod_{i=0}^k \binom{N_i}{r_i},$$

where r_j are the base p digits of r , and we view the right hand side modulo p . In particular, $d_{p^i} = (-1)^{p^i} N_i$ for $i = 0, \dots, k$, where N_i are taken mod p . Thus, we get

$$d_r = (-1)^r \prod_{i=0}^k \binom{(-1)^{p^i} d_{p^i}}{r_i},$$

so $f(d_1, d_p, \dots, d_{p^k}) = 0$ for any $d_1, d_p, \dots, d_{p^k} \in \mathbb{F}_p$ (as N_i are arbitrary). Since the degree of f in each variable is $< p$, this implies that $f = 0$, as desired. \square

Now we are ready to prove (i). Let δ_i be the representative of $(-1)^{p^i} d_{p^i}$ in $[0, p-1]$, and let $d_+ = -\sum_{i \geq 0} \delta_i p^i \in \mathbb{Z}_p$. We have

$$\begin{aligned} \sum_{n=0}^{\infty} d_n z^n &= \prod_{i \geq 0} \sum_{n_i=0}^{p-1} \binom{(-1)^{p^i} d_{p^i}}{n_i} (-z)^{n_i p^i} = \\ &= \prod_{i \geq 0} \sum_{n_i=0}^{p-1} \binom{\delta_i}{n_i} (-z)^{n_i p^i} = \prod_{i \geq 0} (1 - z^{p^i})^{\delta_i} = (1 - z)^{-d_+}. \end{aligned}$$

This proves (i).

(ii) First assume that $p > 2$. Consider the symmetric tensor category $\mathcal{C}' := \text{Supervec} \boxtimes \mathcal{C}$, and define the object $\Pi X := X \otimes \mathbf{k}_- \in \mathcal{C}'$, where \mathbf{k}_- is the odd 1-dimensional supervector space. Then $\Lambda^n X = \mathbb{S}^n \Pi X$ for even n and $\Lambda^n X = \Pi \mathbb{S}^n \Pi X$ for odd n . Thus, (ii) reduces to (i), with $d_-(X) = -d_+(\Pi X)$.

Now consider the case $p = 2$. In this case, we will use an argument analogous to the proof of (i), replacing \mathbb{S}^\bullet by Λ^\bullet (this argument can also be used as an alternative proof for $p > 2$, inserting appropriate signs). Let us describe the necessary changes. First we need to prove an analog of Claim 1:

Claim 3. A' restricts to an endomorphism A_B of $\Lambda^\lambda X$, and the resulting endomorphism is independent on the choice of the b_i .

To see this, note first that $\Lambda^\lambda X \subset \mathbb{S}^\lambda X$, so by Claim 1, the choice of b_i does not matter. To show that A_B lands in $\Lambda^\lambda X$, we need to show that A_B lands in the image of $c_j + 1$ in $X^{\otimes n}$ for each $j \neq \lambda_1, \lambda_1 + \lambda_2, \dots$. Let T_j be the set of all i such that $s_{j,j+1} B_i = B_i$. If $i \notin T_j$, we may choose $b_i \in B_i$ and $b_{i'} \in B_{i'} := s_{j,j+1} B_i$ so that $s_{j,j+1} b_i = b_{i'}$. Thus, it suffices to show that for each $i \in T_j$, b_i maps $\Lambda^\lambda X$ to the image of $c_j + 1$. To this end, note that $s_{j,j+1} b_i = b_i s_{k,l}$, where $s_{k,l} \in S_\lambda$. Since $\Lambda^\lambda X$ is contained in the image of $s_{k,l} + 1$, the image of b_i restricted to $\Lambda^\lambda X$ is contained in the image of $b_i(s_{k,l} + 1) = (s_{j,j+1} + 1)b_i$ on $X^{\otimes n}$. This implies Claim 3.

Now we need to prove an analog of Claim 2.

Claim 4. A' lands in $\Lambda^\lambda X \subset X^{\otimes n}$.

To see this, let U_j be the set of s such that $s_{j,j+1} s = s$. As in Claim 3, it suffices to show that for $s \in U_j$, $c_s g$ restricted to $\Lambda^\mathbf{a} X$ lands in the image of $s_{j,j+1} + 1$. We have $s_{j,j+1} c_s g = c_s g s_{k,l}$, where $s_{k,l} \in S_\mathbf{a}$. We have $\Lambda^\mathbf{a} X \subset \text{Im}(s_{k,l} + 1)$, so the image of the restriction of $c_s g$ to $\Lambda^\mathbf{a} X$ is contained in the image of $c_s g(s_{k,l} + 1) = (s_{j,j+1} + 1)c_s g$. This proves Claim 4.

The rest of the proof is a straightforward generalization of the proof of (i), starting from Lemma 2.5, and the formulas are simpler. Namely,

we have $N = N_0 + N_1p + \dots + N_kp^k$, and the universal polynomial analogous to the polynomial Q_n of Lemma 2.5 has the form

$$Q_n^-(z_0, z_1, \dots) = \prod_{i \geq 0} \binom{z_i}{n_i}.$$

□

Example 2.6. Here are examples of computation of the polynomial P_B constructed in the proof of Lemma 2.4.

1. Let $n = 3$, $\lambda = (2, 1)$. Then $S_n = S_3$ consists of two double cosets of $S_\lambda = S_2 = \langle (12) \rangle$ — the group S_λ itself and the remaining set B of 4 elements: $B = \{(23), (13), (123), (132)\}$. The element of minimal length in B is $g = (23)$ (it has length 1). Then H is the trivial group, and so is S_a . Thus $A_B = (23) + (12)(23) = (23) + (123)$. Hence

$$P_B = \text{Tr}(A_B) = \text{Tr}(A') = \text{Tr}_{X^{\otimes 3}}((23) + (123)) = d_1^2 + d_1.$$

So in characteristic 2 we get $d_3 = d_1d_2 + d_1^2 + d_1$, which gives $d_1d_2 = Q_3(d_1, d_2)$ (as $d_1 \in \mathbb{F}_2$).

2. Let $n = 4$, $\lambda = (3, 1)$. Then $S_n = S_4$ consists of two double cosets of $S_\lambda = S_3 = \langle (12), (23) \rangle$ — the group S_λ itself and the remaining set B of 18 elements. The element of minimal length in B is $g = (34)$ (it has length 1). Thus $H = \langle (12) \rangle = S_a$. Hence $A_B = (34) + (123)(34) + (213)(34) = (34) + (1234) + (2134)$. So

$$\begin{aligned} P_B &= \text{Tr}(A_B) = \text{Tr}(A') = \text{Tr}_{\mathbb{S}^2 X \otimes X \otimes X}((34) + (1234) + (2134)) = \\ &= d_1d_2 + \text{Tr}_{\mathbb{S}^2 X \otimes X \otimes X}((1234) + (234)). \end{aligned}$$

The remaining trace is the trace corresponding to a double coset B' of $S_2 = \langle (12) \rangle$ in S_4 consisting of the elements $(134), (234), (1234), (2134)$ in which the minimal length element is (234) (of length 2). So this trace equals

$$\text{Tr}_{X^{\otimes 4}}((1234) + (234)) = d_1 + d_1^2.$$

Thus we get that $P_B = d_1d_2 + d_1 + d_1^2$. Hence, in characteristic 3 we have

$$d_4 = d_1d_3 + \frac{1}{2}d_1(d_1^2 + d_1) + d_1 + d_1^2 = d_1d_3 - d_1^3 + d_1,$$

which gives $d_1d_3 = Q_3(d_1, d_3)$, as $d_1 \in \mathbb{F}_3$.

Definition 2.7. We call the number $d_+(X^*)$ the *symmetric p -adic dimension of X* , denoted $\text{Dim}_+(X)$, and call the number $d_-(X^*)$ the *exterior p -adic dimension of X* , denoted $\text{Dim}_-(X)$.

Thus, the numbers $\text{Dim}_\pm(X)$ satisfy equations (1), (2).

Example 2.8. Let X be a supervector space. Then $\text{Dim}_+(X) = \text{Dim}_-(X) = \dim(X_0) - \dim(X_1)$, where X_0 and X_1 are the even and odd parts of X .

Let R be the ring of integer-valued polynomials, i.e. polynomials $f \in \mathbb{Q}[t]$ which map integers to integers (or, equivalently, map sufficiently large integers to integers). It is well known that R has a \mathbb{Z} -basis consisting of the binomial coefficients $e_i := \binom{t}{i}$, $i \geq 0$, with multiplication law

$$e_i e_j = \sum_k C_{ij}^k e_k, \quad C_{ij}^k = \frac{k!}{(k-i)!(k-j)!(i+j-k)!}.$$

In particular, the leading coefficient (i.e., that of e_{i+j}) is $\binom{i+j}{i}$.

Recall that if m is not a power of a prime p then there exists an integer $0 < i < m$ such that $\binom{m}{i}$ is not divisible by p . Thus, R is generated by e_j where j is a prime power.

Let $\mathcal{I} \subset \mathbb{Z}[x_1, x_2, \dots]$ be the ideal of polynomials f such that for a symmetric tensor category \mathcal{C} over any field and any $X \in \mathcal{C}$, one has $f(d_1, d_2, \dots) = 0$, where $d_i = \dim \wedge^i X$. Let $\mathcal{R} := \mathbb{Z}[x_1, x_2, \dots]/\mathcal{I}$.

Proposition 2.9. *We have an isomorphism $\psi : \mathcal{R} \rightarrow R$ given by $\psi(x_i) = e_i$.*

Proof. Consider the map $\xi : \mathbb{Z}[x_1, x_2, \dots] \rightarrow R$ sending x_i to e_i . Let $I = \text{Ker} \xi$. We need to show that $I = \mathcal{I}$. Note that I is generated by the elements $f_{ij} := x_i x_j - \sum_k C_{ij}^k x_k$. We claim that $f_{ij} \in \mathcal{I}$, i.e.

$$(4) \quad d_i d_j = \sum_k C_{ij}^k d_k.$$

in any tensor category. To see this, multiply the left hand side by $z^i w^j$ and sum over i, j . We get $(1+z)^d (1+w)^d$, where $d = \text{Dim}_-(X)$. Write this as $(1+z+w+zw)^d$ and expand it as

$$\begin{aligned} (1+z)^d (1+w)^d &= \sum_k d_k (z+w+zw)^k = \\ &= \sum_{i,j,k} \frac{k!}{(k-i)!(k-j)!(i+j-k)!} d_k z^{k-j} w^{k-i} (zw)^{i+j-k} = \\ &= \sum_{i,j,k} \frac{k!}{(k-i)!(k-j)!(i+j-k)!} d_k z^i w^j, \end{aligned}$$

which is exactly the generating function of the right hand sides of (4). Thus, $I \subset \mathcal{I}$, and our job is to establish the opposite inclusion. To do so, let $f \in \mathcal{I}$, and assume that f depends only on x_1, \dots, x_r .

Consider the category $\mathcal{C} = \text{Vect}_{\mathbb{C}}$ of complex vector spaces. Let $X = \mathbb{C}^n$. Then $d_i = \binom{n}{i}$, so $f(1, n, \dots, \binom{n}{r}) = 0$ for any positive integer n . Then $f(1, t, \dots, \binom{t}{r}) = 0$ in R , which implies that $f \in I$, as desired. \square

Remark 2.10. Using Proposition 2.9, we see that the polynomial P_B in Lemma 2.4 is not unique, since R is generated by e_j with j being a prime power (so, d_6 expresses in terms of d_1, \dots, d_5).

Remark 2.11. Let $\tilde{\mathcal{I}} \subset \mathbb{Z}[x_1, x_2, \dots]$ be the ideal of polynomials f such that for a symmetric tensor category \mathcal{C} over any field and any $X \in \mathcal{C}$, one has $f(d_1, d_2, \dots) = 0$, where $d_i = \dim S^i X$. Let $\tilde{\mathcal{R}} := \mathbb{Z}[x_1, x_2, \dots]/\tilde{\mathcal{I}}$. Then similarly to Proposition 2.9 one shows that we have an isomorphism $\tilde{\psi} : \tilde{\mathcal{R}} \rightarrow R$ such that $\tilde{\psi}(x_i) = s_i$, where $s_i := \binom{t+n-1}{n} \in R$.

3. PROPERTIES OF p -ADIC DIMENSIONS

In this section we will study properties of p -adic dimensions Dim_{\pm} . Some of the properties are obvious from the definition — e.g. Dim_{\pm} are invariant under symmetric tensor functors, and $\text{Dim}_{\pm}(X)$ are equal to $\dim(X)$ modulo p . Some less obvious properties are discussed below.

3.1. Values of p -adic dimensions.

Proposition 3.1. $\text{Dim}_{+}(X)$ and $\text{Dim}_{-}(X)$ can take any value in \mathbb{Z}_p .

Proof. This follows from the existence of Deligne’s examples (see [Ha], Subsection 3.3). Namely, e.g. take Deligne’s category $\text{Rep}(S_{\mathcal{U}})$ for an ultrafilter \mathcal{U} , and let V be the tautological object in this category. Then it is easy to see that $\text{Dim}_{+}(V) = \text{Dim}_{-}(V) = t$, where $t = \lim_{\mathcal{U}} n \in \mathbb{Z}_p$, and t can be arbitrary. \square

3.2. Additivity on exact sequences. If Y is a \mathbb{Z}_{+} -filtered object of a symmetric tensor category \mathcal{C} , then we will denote by $\text{gr}Y$ the associated graded object $\bigoplus_i F_{i+1}Y/F_iY$; so, $\dim Y = \dim \text{gr}Y$. Also note that if Y is \mathbb{Z}_{+} -filtered then the symmetric algebra SY carries an induced filtration.

Let

$$0 \rightarrow X \rightarrow Y \rightarrow Z \rightarrow 0$$

be a short exact sequence; it induces a 2-step filtration on Y .

Proposition 3.2. *Suppose that the natural surjective morphism*

$$\phi_{+} : S(X \oplus Z) = S(\text{gr}Y) \rightarrow \text{gr}SY$$

is an isomorphism. Then $\text{Dim}_+(Y) = \text{Dim}_+(X) + \text{Dim}_+(Z)$. Likewise, if the natural surjective morphism

$$\phi_- : \wedge(X \oplus Z) = \wedge(\text{gr}Y) \rightarrow \text{gr} \wedge Y$$

is an isomorphism then $\text{Dim}_-(Y) = \text{Dim}_-(X) + \text{Dim}_-(Z)$.

Proof. For a graded ind-object $W = \bigoplus_{n \geq 0} W_n$ of \mathcal{C} , define its Hilbert series by

$$h_W(z) = \sum_{n \geq 0} \dim W_n z^n.$$

Then by definition of p -adic dimensions, for any object $V \in \mathcal{C}$, $h_{S^p V}(z) = (1-z)^{-\text{Dim}_+(V)}$ and $h_{\wedge^p V}(z) = (1+z)^{\text{Dim}_-(V)}$. Thus, the result is obtained by computing the Hilbert series of SY as the product of Hilbert series of SX and SZ , using that $\text{gr}SY = S(X \oplus Z) = SX \otimes SZ$, and similarly for exterior algebras. \square

Example 3.3. The assumption in Proposition 3.2 cannot be removed, and, in fact, its conclusion can fail at least in characteristic 2. To give an example, let D be the Hopf algebra $\mathbf{k}[d]/d^2$ with d being a primitive element and triangular structure $R = 1 \otimes 1 + d \otimes d$. Consider the non-semisimple symmetric tensor category $\text{Rep}D$. This category plays the role of the category of supervector spaces in characteristic 2. Take $Y = D$, and $X = Z = \mathbf{1}$. Then the map ϕ_+ is not an isomorphism; in fact, $S^n D = D$ for $n \geq 1$ (see [V], Subection 1.5). Thus, since $\dim(D) = 0$, we get that $\text{Dim}_+(Y) = 0$. Likewise, $\wedge^n D = D$ for $n \geq 1$, so $\text{Dim}_-(Y) = 0$. On the other hand, $\text{Dim}_\pm(X) = \text{Dim}_\pm(Z) = 1$, so $\text{Dim}_\pm(Y) \neq \text{Dim}_\pm(X) + \text{Dim}_\pm(Z)$.

Remark 3.4. The symmetric category $\text{Rep}D$ appeared 50 years ago in algebraic topology. Namely, it is explained in [ATo], Corollary 7.8 that the K-theory of a topological space with coefficients in $\mathbb{Z}/2$ is a $\mathbb{Z}/2$ -graded commutative algebra in $\text{Rep}D$, with d of degree 1 (the Bockstein homomorphism), and in fact it does not admit (universal) multiplications which are commutative in the usual sense (see also [AY], (2.17)).

Question 3.5. Suppose $p > 2$, and Y has a finite filtration. Are the natural epimorphisms $\phi_+ : S(\text{gr}Y) \rightarrow \text{gr}SY$, $\phi_- : \wedge(\text{gr}Y) \rightarrow \text{gr} \wedge Y$ isomorphisms?

Clearly, for this question it suffices to consider 2-step filtrations. By Proposition 3.2, a positive answer to Question 3.5 would imply that Dim_\pm are additive on exact sequences.

Example 3.3 shows that the answer is “no” for $p = 2$. However, the following proposition shows that this type of counterexamples does not exist for $p > 2$.

Proposition 3.6. *Let H be a cotriangular connected Hopf algebra over a field \mathbf{k} of characteristic $p > 2$ (i.e., every simple H -comodule is trivial, and the category of comodules is symmetric). Then for any \mathbb{Z}_+ -filtered finite dimensional H -comodule Y , the natural surjections ϕ_+, ϕ_- are isomorphisms.*

Proof. Let us prove the statement for ϕ_+ ; for ϕ_- the proof is similar.

First of all, it suffices to prove the statement for any refinement of the filtration on Y . In particular, we may take a maximal refinement, in which the successive quotients are trivial 1-dimensional modules, i.e., $Y_0 := \text{gr}Y$ is a vector space with a trivial action of H .

Let $\text{Rees}(Y) := \prod_{j \geq 0} \hbar^j F_j Y$. Then $S(\text{Rees}(Y))$ is a formal deformation of $SY_0 = \mathbf{k}[x_1, \dots, x_n]$ as a quadratic algebra. Our job is to show that this deformation is flat. By Drinfeld’s “Koszul deformation principle” ([PP], Introduction, p. ix), it suffices to check this in degrees 2, 3. If $p > 3$, then the action of S_n on $Y_0^{\otimes n}$ is the same as in $Y^{\otimes n}$ for $n = 2, 3$ (as representations of S_n are semisimple and hence deformation-theoretically rigid), so the condition is clearly satisfied, and we get flatness, as desired.

Consider now the case $p = 3$. Then the same argument works in degree 2, and it remains to analyze degree 3.

We claim that the deformation of the S_3 -action from $Y_0^{\otimes 3}$ to $Y^{\otimes 3}$ is still trivial. To show this, it suffices to check that $H^1(S_3, \text{End}(Y_0^{\otimes 3})) = 0$. For this, it is enough to show that for any vector space W , we have $H^1(S_3, W^{\otimes 3}) = 0$ (then we can take $W = \text{End}(Y_0)$). To prove this, let w_i be a basis of W . Using this basis, we can decompose $W^{\otimes 3}$ as an S_3 -module into three kinds of modules:

1) The span of the orbit of $w_i \otimes w_j \otimes w_m$ for i, j, m distinct: the regular representation of S_3 . This gives the zero cohomology by the Shapiro lemma.

2) The span of the orbit of $w_i \otimes w_i \otimes w_j$ for i, j distinct: the induced module from the trivial module over S_2 . So again by the Shapiro lemma we get that the cohomology in question is $H^1(S_2, \mathbf{k}) = 0$ (as $p = 3$).

3) The span of $w_i \otimes w_i \otimes w_i$. This gives the cohomology $H^1(S_3, \mathbf{k}) = \text{Hom}(S_3, \mathbf{k}) = 0$, as $\text{char}(\mathbf{k}) = 3$. We are done. \square

Remark 3.7. 1. Note that this does not work in characteristic 2, even in degree 2, since for $\text{char}(\mathbf{k}) = 2$, $H^1(S_2, \mathbf{k}) = \mathbf{k}$ is not zero.

2. This proof extends verbatim to the case when simple H -comodules factor through a commutative Hopf subalgebra $H_0 \subset H$ (with trivial

triangular structure), i.e. $H_0 = \mathcal{O}(G)$, where G is an affine group scheme.

3. We can extend this further to the setting as above, except that H is a Hopf superalgebra, G is an affine supergroup scheme, and $\mathcal{O}(G)$ has a triangular structure defined by some central element $u \in G$ of order 2. In this case, the proof is the same, except we have to show for $p = 3$ that $H^1(S_3, \text{End}_{\text{even}}(Y_0^{\otimes 3})) = 0$, where the action of S_3 on $\text{End}_{\text{even}}(Y_0^{\otimes 3})$ takes into account the signs. So case (1) above is the same, in case (3) we must have all w_i even, so we get $H^1(S_3, \mathbf{k}) = 0$, and in case (2) we get two subcases:

(2a) w_i is even; then we get the induced module from the trivial S_2 -module, so we get $H^1(S_2, \mathbf{k}) = 0$; and

(2b) w_i is odd; then we get the induced module from the sign character \mathbf{k}_- of S_2 , and we get $H^1(S_2, \mathbf{k}_-)$, which is again zero.

3.3. Compatibility with the λ -ring structure. Recall now that a pre- λ -ring is a commutative unital ring with additional operations λ^i satisfying the following axioms (see e.g. [AT, Y]).

- (1) $\lambda^0(x) = 1$
- (2) $\lambda^1(x) = x$
- (3) $\lambda^n(1) = 0$ for $n > 1$
- (4) $\lambda^n(x + y) = \sum_{i+j=n} \lambda^i(x)\lambda^j(y)$

It is an easy exercise to verify that in a symmetric tensor category \mathcal{C} , if for every short exact sequence $0 \rightarrow X \rightarrow Y \rightarrow Z \rightarrow 0$ the natural surjective map $\psi : \wedge(X \oplus Z) = \wedge(\text{gr}Y) \rightarrow \text{gr} \wedge Y$ is an isomorphism (as in Proposition 3.2) then the Grothendieck ring $\text{Gr}(\mathcal{C})$ inherits the structure of a pre- λ -ring with the structure maps $\lambda^i([X]) = [\wedge^i X]$.

A pre- λ -ring is a λ -ring if it satisfies two additional (families of) axioms encoding plethysm:

- (5) $\lambda^n(xy) = P_n(\lambda^1(x), \lambda^2(x), \dots, \lambda^n(x), \lambda^1(y), \lambda^2(y), \dots, \lambda^n(y))$
- (6) $\lambda^m(\lambda^n(x)) = P_{m,n}(\lambda^1(x), \lambda^2(x), \dots, \lambda^{mn}(x))$

where P_n and $P_{n,m}$ are certain universal polynomials with integer coefficients (see [AT, Y]).

For instance, \mathbb{Z} and \mathbb{Z}_p are λ -rings, with $\lambda^i(x) = \binom{x}{i}$. Also, the Grothendieck ring of any symmetric tensor category over a field of characteristic zero is a λ -ring, with $\lambda^i = \wedge^i$ (this follows from the fact that this holds for the category of Schur functors, i.e. that the ring Λ of symmetric functions over \mathbb{Z} is a λ -ring).

Proposition 3.8. *Suppose that the Grothendieck ring $\text{Gr}(\mathcal{C})$ is a λ -ring, with $\lambda^i([X]) = [\wedge^i X]$ for all $X \in \mathcal{C}$. Then $\text{Dim}_- : \text{Gr}(\mathcal{C}) \rightarrow$*

\mathbb{Z}_p is a homomorphism of λ -rings. In particular, $\text{Dim}_-(X \otimes Y) = \text{Dim}_-(X)\text{Dim}_-(Y)$ and $\text{Dim}_-(\wedge^i X) = \binom{\text{Dim}_-(X)}{i}$.

Proof. Recall that the ring $W(\mathbb{F}_p)$ of big Witt vectors of \mathbb{F}_p is isomorphic to $1 + z\mathbb{F}_p[[z]]$ as an abelian group, and has a λ -ring structure with the following adjunction property: if R is a λ -ring and $f : R \rightarrow \mathbb{F}_p$ is a map of rings, then the map $F := 1 + \sum_{j>0} (f \circ \lambda^j) z^j$ is a map of λ -rings from R to $W(\mathbb{F}_p)$, which, when composed with the ‘‘coefficient of z ’’ map of rings from $W(\mathbb{F}_p)$ to \mathbb{F}_p , gives f .

By Lemma 2.2 we may think of the usual categorical dimension as a map of rings $\text{dim} : \text{Gr}(\mathcal{C}) \rightarrow \mathbb{F}_p$. Applying the above adjunction to this gives a map of λ -rings $\text{Dim} : \text{Gr}(\mathcal{C}) \rightarrow W(\mathbb{F}_p)$. Explicitly this map sends an object X to $\sum_j \text{dim}(\wedge^j X) z^j = (1 + z)^{\text{Dim}_-(X)}$ by Theorem 2.3.

The copy of \mathbb{Z}_p in $W(\mathbb{F}_p)$ given by the inclusion $\alpha \rightarrow (1 + z)^\alpha$, was characterized by Elliot ([El] Proposition 9.3) as the largest sub- λ -ring of $W(\mathbb{F}_p)$ for which the Adams operations act by the identity, which implies that the λ -operations are given by the usual binomial formulas there. We see that Dim lands inside this sub- λ -ring and coincides with Dim_- under its identification with \mathbb{Z}_p .

In particular this implies that Dim_- is a map of λ -rings from $\text{Gr}(\mathcal{C})$ to \mathbb{Z}_p (equipped with the binomial λ -ring structure) and hence satisfies $\text{Dim}_-(X \otimes Y) = \text{Dim}_-(X)\text{Dim}_-(Y)$ and $\text{Dim}_-(\wedge^i X) = \binom{\text{Dim}_-(X)}{i}$, as desired. □

This proposition applies to classical categories (supergroup representations, etc.) as well as to $\text{Rep}(S_U)$, $\text{Rep}(GL_U)$. However, it is possible for both the assumption and conclusion of Proposition 3.8 to fail in any characteristic $p \geq 5$.

To give an example, consider the Verlinde category $\mathcal{C} = \text{Ver}_p$, see [O] and references therein. This is the quotient of the category of representations of $\mathbb{Z}/p\mathbb{Z}$ by negligible morphisms. In Ver_p we have an object L_j coming from the j -dimensional Jordan block representation of $\mathbb{Z}/p\mathbb{Z}$ ($1 \leq j \leq p-1$). Let $V = L_2$. Then L_2 satisfies the equation $P_{p-1}(L_2) = 0$, where P_n is the n -th ultraspherical polynomial defined by the formula $P_n(q + q^{-1}) = \sum_{i=0}^n q^{n-2i}$. We claim that P_{p-1} has no roots in \mathbb{Q}_p . Indeed, the roots of P_p are of the form $\zeta + \zeta^{-1}$, where ζ is a root of unity of order $2p$. If such a root were in \mathbb{Q}_p then ζ would have been in a quadratic extension of \mathbb{Q}_p ; but $\mathbb{Q}_p(\zeta^2)$ is well known to be a ramified extension of degree $p-1 > 2$. This shows that there can't even be a ring homomorphic lift of the dimension map to \mathbb{Z}_p . In

particular, Dim_- fails to be multiplicative in this example, and hence $\text{Gr}(\mathcal{C})$ is not (naturally) a λ -ring.

Example 3.9. Let $p = 5$. Then $X = L_3$ satisfies the quadratic equation $X^2 - X - 1 = 0$ in $\text{Gr}(\mathcal{C})$. This equation has no roots in \mathbb{Z}_5 , so there is no homomorphisms from $\text{Gr}(\mathcal{C})$ to \mathbb{Z}_5 .

3.4. The Koszul complex and equality of p -adic dimensions.

For supervector spaces both Dim_+ and Dim_- recover the usual notion of dimension of a supervector space, and in particular we have

$$(5) \quad \text{Dim}_+(X) = \text{Dim}_+(X^*) = \text{Dim}_-(X) = \text{Dim}_-(X^*).$$

Therefore, we also get these equalities for any symmetric tensor category admitting a fiber functor to supervector spaces (i.e. supergroup representations) as well as interpolations of such categories (i.e., Deligne categories $\text{Rep}(S_U)$, $\text{Rep}(GL_U)$). However, in a general symmetric tensor category, (5) can fail in any characteristic $p \geq 5$.

To give an example, consider again the category Ver_p . It is easy to see that $\wedge^2 V = L_1 = \mathbf{1}$ and $\wedge^i V = 0$ for all $i > 2$. Also, $\dim(L_j) = j$. Thus, $\text{Dim}_-(V) = 2$. At the same time, $SV = L_1 \oplus L_2 \oplus \dots \oplus L_{p-1}$. Hence,

$$\sum_{j=0}^{\infty} \dim(S^j V) z^j = 1 + 2z + \dots + (p-1)z^{p-2} = (1-z)^{-2}(1-z^p).$$

Thus, $\text{Dim}_+(V) = 2 - p$, and $\text{Dim}_+(V) \neq \text{Dim}_-(V)$.

On the other hand, in Ver_p we still have $\text{Dim}_+(X) = \text{Dim}_+(X^*)$ and $\text{Dim}_-(X) = \text{Dim}_-(X^*)$. In fact, we don't know if these equalities can ever fail; see Question 3.13 below.

It is therefore an interesting question when all or some of the equalities (5) hold. The following two propositions give sufficient conditions for this.

To state the first proposition, recall that to any object $X \in \mathcal{C}$ we can attach its *Koszul complex* $K^\bullet(X) = SX \otimes \Lambda^\bullet X$, with the usual Koszul differential $\partial : K^i(X) \rightarrow K^{i-1}(X)$. Namely, we have the multiplication morphism $X^* \otimes \wedge^{i-1} X^* \rightarrow \wedge^i X^*$, which after dualization defines the morphism $\mu : X^* \otimes \Lambda^i X \rightarrow \Lambda^{i-1} X$. Then the differential ∂ is given by the formula

$$\partial = (m \otimes \mu) \circ P_{23} \circ (\text{coev}_X \otimes 1_{SX} \otimes 1_{\Lambda X}),$$

where $\text{coev}_X : \mathbf{1} \rightarrow X \otimes X^*$ is the coevaluation, P_{23} is the permutation of the second and the third factor, and $m : X \otimes SX \rightarrow SX$ is the multiplication.

The Koszul complex is a \mathbb{Z}_+ -graded complex under diagonal grading. By a classical result of commutative algebra, in the category of vector spaces, this complex is exact. More precisely, it is exact in positive homological degrees and has homology equal to $\mathbf{1}$ in degree 0; i.e., it is a resolution of the augmentation module by free SX -modules. Note also that for vector spaces $\Lambda X = \wedge X$, so the Koszul complex may be defined as $K^\bullet(X) = SX \otimes \wedge^\bullet X$, the differential graded algebra generated by the superspace $X \oplus \Pi X$ with differential $d(x, \xi) = (0, x)$ (where Π is the change of parity functor).

Also, if $i + j = n$ then

$$S^i \Pi X \otimes \Lambda^j \Pi X = \Pi^n (\wedge^i X \otimes S^j X) = \Pi^n (S^j X^* \otimes \wedge^i X^*)^*.$$

Thus, the homogeneous components of $K^\bullet(X)$ are dual to those of $K^\bullet(\Pi X^*)$ up to parity change. This implies that the Koszul complex is exact in the category Supervec , and hence for representation categories of supergroups and their interpolations (Deligne categories).

Also, it is easy to show that in any symmetric tensor category over \mathbf{k} , the Koszul complex is exact if $\text{char}(\mathbf{k}) = 0$ (as it is exact in the category of Schur functors), and is exact in diagonal degrees $< p$ for characteristic p . However, the following proposition implies that in general, the complex $K^\bullet(X)$ can fail to be exact in any characteristic $p \geq 5$.

Proposition 3.10. *$\text{Dim}_-(X^*) = \text{Dim}_+(X)$ if and only if the Euler characteristic of the Koszul complex $K^\bullet(X)$ is zero outside of diagonal degree zero.*

Proof. It is immediate from the definition of Dim_\pm that $\text{Dim}_-(X^*) = \text{Dim}_+(X)$ iff

$$\left(\sum_{j \geq 0} (-1)^j \dim(S^j X) z^j \right) \left(\sum_{j \geq 0} \dim(\Lambda^j X) z^j \right) = 1$$

Taking the coefficient of z^n for $n > 0$ tells us that

$$\sum_j (-1)^j \dim(S^j X) \dim(\Lambda^{n-j} X) = 0$$

and one can easily recognize the left hand side as being the Euler characteristic of the degree n part of the Koszul complex. \square

Next, we will give a sufficient condition for the equality $\text{Dim}_+(X) = \text{Dim}_-(X)$ which is related to Proposition 3.8.

Proposition 3.11. *Suppose $\text{char}(\mathbf{k}) > 2$, \mathcal{C} contains Supervec , and Dim_- is multiplicative on \mathcal{C} (e.g., the Grothendieck ring $\text{Gr}(\mathcal{C})$ is a*

λ -ring with the usual operations). Then $\text{Dim}_-(X) = \text{Dim}_+(X)$ for all $X \in \mathcal{C}$.

Proof. Let \mathbf{k}_- be the nontrivial invertible object in $\text{Supervec} \subset \mathcal{C}$. We have that $\text{Dim}_-(\mathbf{k}_-) = \text{Dim}_+(\mathbf{k}_-) = -1$, and moreover \mathbf{k}_- exchanges the roles of symmetric and exterior powers in the following sense:

$$\wedge^j(\mathbf{k}_- \otimes X) = \mathbf{k}_-^{\otimes j} \otimes S^j X$$

Now, since Dim_- is multiplicative, we have:

$$\begin{aligned} (1+z)^{-\text{Dim}_-(X)} &= (1+z)^{\text{Dim}_-(\mathbf{k}_- \otimes X)} \\ &= \sum_j \dim(\wedge^j(\mathbf{k}_- \otimes X)) z^j \\ &= \sum_j \dim(\mathbf{k}_-^{\otimes j} \otimes S^j X) z^j \\ &= \sum_j (-1)^j \dim(S^j X) z^j \\ &= (1 - (-z))^{-\text{Dim}_+(X)} = (1+z)^{-\text{Dim}_+(X)} \end{aligned}$$

In particular we see that $\text{Dim}_-(X) = \text{Dim}_+(X)$, as desired. \square

Remark 3.12. If \mathcal{D} is any tensor category over \mathbf{k} then Proposition 3.11 may be applied to the category $\mathcal{C} := \text{Supervec} \boxtimes \mathcal{D}$, provided that Dim_- is multiplicative on this category.

3.5. Categorical Brauer characters. One can use p -adic dimension to define Brauer characters in the categorical setting. Let G be a finite group acting on $X \in \mathcal{C}$. Then for each $g \in G$ and each root of unity $\zeta \in \mathbf{k}$, we have the generalized eigenobject $X(g, \zeta)$, the direct limit of kernels of $(g - \zeta)^N|_X$ as $N \rightarrow \infty$. Let K be the maximal unramified extension of \mathbb{Q}_p (obtained by adding all roots of unity of orders prime to p). Then by Hensel's lemma, ζ has a canonical lift $\hat{\zeta}$ to K . Define

$$\chi_X(g) = \sum_{\zeta} \text{Dim}_+(X(\zeta, g)) \hat{\zeta} \in K.$$

We may call this the Brauer character of G on X . Note that if X is a usual representation then it is the usual Brauer character. One can also make a similar definition using Dim_- .

3.6. Questions. Here are some questions which could be a subject of further research. Let \mathcal{C} be a symmetric tensor category over a field \mathbf{k} of characteristic $p > 0$. Let $X \in \mathcal{C}$.

Question 3.13. 1. Do $(S^i X)^*$ and $S^i X^*$ always define the same class in the Grothendieck group $\text{Gr}(\mathcal{C})$? Same question for $(\wedge^i X)^*$, $\wedge^i X^*$.

2. Is it true that $\dim(S^i X) = \dim(S^i X^*)$, $\dim(\wedge^i X) = \dim(\wedge^i X^*)$? In other words, is it true that $\text{Dim}_+(X) = \text{Dim}_+(X^*)$, $\text{Dim}_-(X) = \text{Dim}_-(X^*)$?

A positive answer to (1) would imply one to (2). Also, (1) holds for representation categories of finite groups, as the objects in question have the same Brauer characters (using Molien's formula).

Question 3.14. Suppose $p = 2$ or $p = 3$.

1. Is it true that $\text{Dim}_+ = \text{Dim}_-$?

2. If the maps ϕ_+ , ϕ_- of Proposition 3.2 are always bijective for \mathcal{C} , is it true that Dim_\pm are ring homomorphisms $\text{Gr}(\mathcal{C}) \rightarrow \mathbb{Z}_p$? is $\text{Gr}(\mathcal{C})$ a λ -ring with the usual exterior power operations in this case?

3. Is the Koszul complex $K^\bullet(X)$ always exact?

Question 3.15. Suppose for some $C \geq 1$ one has $\text{length}(X^{\otimes n}) \leq C^n$ for any $n \geq 1$. Does it follow that $\text{Dim}_\pm(X)$ are integers?

For the next question, let λ be a partition of n , and π_λ be the corresponding Specht module over $\mathbf{k}S_n$. Define the Schur functor $\mathcal{S}^\lambda X := \pi_\lambda \otimes_{S_n} X^{\otimes n}$. E.g., $\mathcal{S}^{(n)} X = S^n X$ and $\mathcal{S}^{(1^n)} X = \wedge^n X$. Recall that in characteristic zero, there exists a polynomial P_λ such that $\dim \mathcal{S}^\lambda X = P_\lambda(\dim X)$. However, as we have seen, in positive characteristic this fails even for the functors S^i and \wedge^i , and the situation is more complicated. This motivates the following question.

Question 3.16. Suppose $\text{Dim}_+ = \text{Dim}_-$, and $\text{Dim}_+(X) = \text{Dim}_-(X) = t$. When can one express $\dim \mathcal{S}^\lambda X$ in terms of t ?

Note that there are other versions of Schur functors: we can replace the functor $\pi_\lambda \otimes_{S_n} ?$ with the functor $\text{Hom}_{S_n}(\pi_\lambda, ?)$ (which for $\lambda = (n)$ and (1^n) yields the functors \mathbb{S} and Λ , respectively), and can also replace π_λ with π_λ^* . One may also consider the corresponding derived functors. The behavior of dimensions under such functors is an interesting open problem.

REFERENCES

- [AT] M. F. Atiyah, D. O. Tall, Group representations, λ -rings and the J-homomorphism, *Topology* 8: 253-297, 1969.
- [ATo] S. Araki, H. Toda, Multiplicative structures in mod q cohomology theories, II, *Osaka J. Math.* 3, 1966, 81-120.
- [AY] S. Araki, Z. Yosimura, Differential Hopf algebras modelled on K-theory mod p , I, *Osaka J. Math.* 8 (1971), 151-206.

- [De1] P. Deligne, Categories Tannakiennes, In The Grothendick Festschrift, Vol. II, Prog. Math. 87 (1990), p.111-195.
- [De2] P. Deligne, La Catégorie des Représentations du Groupe Symétrique S_t , lorsque t n'est pas un Entier Naturel, Algebraic Groups and Homogeneous Spaces, Mumbai 2004. (TIFR 2007) pp. 209-273.
- [De3] P. Deligne, Letter to Ostrik, March 4, 2015.
- [DM] P. Deligne and J.S. Milne, Tannakian Categories. Hodge cycles, motives, and Shimura varieties. Lecture Notes in Mathematics, 900. Springer, 1982.
- [El] J. Elliot, Binomial rings, integer-valued polynomials, and λ -rings, Journal of Pure and Applied Algebra, 207, 2006, p.165-185
- [EGNO] P. Etingof, S. Gelaki, D. Nikshych, V. Ostrik, Tensor categories, AMS, Providence, 2015.
- [Ha] N. Harman, Stability and periodicity in the modular representation theory of symmetric groups, arXiv:1509.06414.
- [O] V. Ostrik, On symmetric fusion categories in positive characteristic, arXiv:1503.01492.
- [PP] A. Polishchuk, L. Positselski, Quadratic Algebras. University Lecture Series 37, American Mathematical Society, Providence, RI, 2005.
- [S] L. Solomon, A Mackey formula in the group ring of a Coxeter group, J. of Algebra, v.41, 1976, pp.255-268.
- [St] R. Stanley, Enumerative Combinatorics, v.2, Cambridge University Press, 1999.
- [V] S. Venkatesh, Hilbert Basis Theorem and Finite Generation of Invariants in Symmetric Fusion Categories in Positive Characteristic, arXiv:1507.05142.
- [Y] D. Yau, Lambda-rings, World Scientific, 2010.

ETINGOF: DEPARTMENT OF MATHEMATICS, MASSACHUSETTS INSTITUTE OF TECHNOLOGY, CAMBRIDGE, MA 02139, USA

E-mail address: `etingof@math.mit.edu`

HARMAN: DEPARTMENT OF MATHEMATICS, MASSACHUSETTS INSTITUTE OF TECHNOLOGY, CAMBRIDGE, MA 02139, USA

E-mail address: `nharman@mit.edu`

OSTRIK: DEPARTMENT OF MATHEMATICS UNIVERSITY OF OREGON EUGENE, OR 97403, USA

E-mail address: `vostrik@math.uoregon.edu`